

Information and Consent Form for Fat-Dissolving Injection “BNLS”

✿ About BNLS

BNLS is a fat-dissolving injection that combines Sodium Deoxycholate—a newly added active ingredient—with the original plant-derived components.

Note: Sodium Deoxycholate is a medically approved ingredient recognized by the U.S. FDA in recent years for its fat-dissolving effects. It acts by disrupting fat cell membranes and breaking down triglycerides, effectively dissolving fat cells.

✿ Expected Effects

BNLS promotes the breakdown of fat, skin tightening, and improved lymphatic circulation. These three effects work together to improve blood and lymph flow, helping to eliminate dissolved fat and waste products accumulated in the body.

✿ Applicable Treatment Areas

BNLS can be used on the following areas:

- Forehead •Nose and sides of the nose •Cheekbones
- Cheeks (including around the nasolabial folds) •Lower cheeks and sides of the face
- Under the chin (submental area) •Upper arms •Thighs •Abdomen •Back

✿ Precautions

•Swelling and Bruising

Mild pain, swelling, or bruising (subcutaneous bleeding) may occur after the injection. Swelling typically subsides within 2 to 3 days, while bruising gradually fades over 3 to 4 weeks. In rare cases, pigmentation may occur, but this usually improves gradually over several months. To minimize the risk of infection, please ensure the injection site remains clean and dry.

•While individual results may vary, an average of 3 to 4 treatment sessions is generally recommended.

•As inflammation, swelling, or redness may temporarily worsen in the days following the procedure, please avoid strenuous exercise, sauna use, and alcohol consumption.

•If redness or swelling persists, please consult with our clinic’s physician as soon as possible.

•Vasovagal reactions may occur either before or after the injection, especially in patients experiencing high levels of anxiety or psychological stress. Overstimulation of the nerves can cause a sudden drop in blood pressure, resulting in dizziness, nausea, or even loss of consciousness. If you have experienced such symptoms in the past, please inform our staff in advance.

•Although the injection is administered under sterile conditions after disinfecting the skin, there is a rare risk of infection. To minimize this risk, please keep the injection site clean.

•In certain cases, we may ask for your cooperation in undergoing a blood test to help prevent infection due to needlestick injuries or other medical incidents.

❖ Patients Who Cannot Receive Treatment or Require Special Precaution

You may not be eligible for treatment if any of the following apply:

- You are pregnant, breastfeeding, or may possibly be pregnant
- You have microvascular disorders or vascular insufficiency
- You have an immunodeficiency or autoimmune disease
- You are taking anticoagulant ("blood thinning") medication
- You are currently undergoing chemotherapy
- You have ulcers, eczema, psoriasis, or infections on the treatment area

Please be sure to follow all instructions provided by your treating physician throughout the treatment period. Failure to do so may result not only in reduced treatment effectiveness but also in potential health complications.

If you experience any unusual symptoms during the treatment process, please promptly report them to a doctor or nurse, and follow the physician's guidance after a proper medical assessment.

At our clinic, we take photographs for the purpose of documenting treatment progress in your medical records. If you do not consent to these photographs, it may not be possible to adequately monitor your progress, and we may not be able to appropriately address inquiries about treatment results or manage complications (e.g., skin reactions or nerve symptoms) that may occur.

Please be aware that additional treatment fees may apply in the event of complications.